

AD-A031 946

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER  
ON THE RANGE OF ACCRETIVE OPERATORS.(U)  
SEP 76 M G CRANDALL; A PAZY

F/G 12/1

UNCLASSIFIED

MRC-TSR-1682

DAA029-75-C-0024

NL

1 OF 1  
AD  
A031946





ADA031946

MRC Technical Summary Report #1682

ON THE RANGE OF ACCRETIVE OPERATORS

Michael G. Crandall & Amnon Pazy

**Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706**

September 1976

Received August 20, 1976

*See Form 1473*

**Approved for public release  
Distribution unlimited**

Sponsored by

U. S. Army Research Office  
P.O. Box 12211  
Research Triangle Park  
North Carolina 27709

and

National Science Foundation  
Washington, D. C. 20550

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

ON THE RANGE OF ACCRETIVE OPERATORS

Michael G. Crandall<sup>†</sup> & Amnon Pazy

Technical Summary Report #1682  
September 1976

ABSTRACT

The solvability of the nonlinear operator equation  
 $w = x + Bx$ , where  $B$  is accretive in a general Banach  
space  $X$  is studied by means of discrete approximations.  
In particular, an algorithm is given for solving the equation  
in the case that  $B$  is continuous and everywhere defined.

ADDITIONAL FOR	
NTIS	White Section <input checked="" type="checkbox"/>
F C	Black Section <input type="checkbox"/>
UNCLASSIFIED	JUSTIFICATION <input type="checkbox"/>
BY	
DISTRIBUTION/AVAILABILITY CODES	
Div.	ANAL. REPT. OR SPECIAL
A	

AMS(MOS) Classification No. 47H15

Key Words: Accretive operator, iteration scheme, range.

Work Unit No. 1 - Applied Analysis

<sup>†</sup> Supported in part by NSF grant MCS 76-10227



## ON THE RANGE OF ACCRETIVE OPERATORS

Michael G. Crandall<sup>†</sup> & Amnon Pazy

### Introduction.

Let  $X$  be a real Banach space and let  $B : D(B) \subseteq X \rightarrow X$  be a nonlinear operator.  $B$  is called accretive if

$$\|x_1 - x_2\| \leq \|x_1 - x_2 + \lambda(Bx_1 - Bx_2)\| \text{ for } \lambda > 0, x_1, x_2 \in D(B).$$

If  $B$  is accretive, then  $B$  is m-accretive if  $X = R(I + B)$ , i.e. for every  $w \in X$  there is an  $x \in D(B)$  such that  $w = x + Bx$ . One of the first results in the study of accretive operators was obtained by G. Minty [5] and implied that every continuous everywhere defined accretive operator in a Hilbert space is m-accretive. This latter result was extended to general Banach spaces by R. Martin [4]. The known proofs of Martin's theorem employ the solvability of the initial-value problem

$$(1) \quad \begin{cases} \frac{du}{dt} + Au = 0 \\ u(0) = x \end{cases}$$

where  $A$  is continuous and accretive on  $X$ . The existence theory for equation (1) has been generalized to allow cases in which  $A$  is neither continuous nor single-valued. For recent developments see, e.g., Y. Kobayashi [3] and M. Pierre [7,8]. This existence theory is rather technical and complex. The present paper was motivated by the desire to find a proof of the above mentioned theorem of Martin which is direct, constructive and which does not rely on the solvability of (1).

---

<sup>†</sup>Supported in part by NSF grant MCS 76-10227

If  $B$  is continuous, everywhere defined and accretive we show in Theorem 2 of Section 1 how to choose numbers  $\alpha_k \in (0,1]$  so that the sequence defined by  $x_{k+1} = \alpha_{k+1}x_k - (1-\alpha_{k+1})(Bx_k - z)$  converges to the unique solution  $x_\infty$  of  $x_\infty + Bx_\infty = z$ . (The choice of  $\alpha_{k+1}$  depends on  $x_k$  and  $\alpha_1 + \dots + \alpha_k$ .) In fact, our main results easily adapt to provide elementary proofs (i.e., proofs not relying on (1)) of the strong generalizations of Martin's theorem obtained in [3] and [6]. In particular, the perturbation theorem of Webb [9] as generalized by Barbu [1] follows easily.

The main results are stated in Section 1 and proved in Section 2. Then various known results are obtained as applications in the final Section 3.

#### 1. The Main Results.

Let  $B$  be a mapping from  $X$  to the subsets of  $X$  which is accretive, i.e.

$$(1.1) \quad \|x_1 - x_2 + \lambda(y_1 - y_2)\| \geq \|x_1 - x_2\|$$

for  $\lambda > 0$ ,  $y_1 \in Bx_1$ ,  $x_1 \in D(B) = \{x \in X : Bx \neq \emptyset\}$ . Given  $w \in X$  we consider the solvability of the problem  $w \in R(I + B)$ , i.e. can we find  $x \in X$  such that  $w \in x + Bx$  (equivalently,  $w - x \in Bx$ ). Replacing  $B$  by  $B_w$  where  $B_w x = Bx - w$  we reduce to the case  $w = 0$ .

Definition: A sequence  $\{x_k\}_{k=0}^\infty$  is admissible for the problem  $0 \in R(I + B)$  if there exist  $y_k \in Bx_k$  and numbers  $h_k > 0$ ,  $k = 1, 2, \dots$  such that

$$(1.2) \quad \begin{cases} (i) & \sum_1^\infty h_k = \infty \\ (ii) & \sum_1^\infty \|x_{k+1} - x_k + h_{k+1}(x_{k+1} + y_{k+1})\| < \infty. \end{cases}$$



The first result is:

Theorem 1. Let  $B$  be accretive and  $\{x_k\}$  be an admissible sequence for  $0 \in R(I + B)$ . Then

$$(a) \quad x_\infty = \lim_{k \rightarrow \infty} x_k \text{ exists.}$$

Moreover

$$(b) \quad (1+\lambda) \|x - x_\infty\| \leq \|x - x_\infty + \lambda(y+x)\| \quad \text{for } \lambda > 0, x \in D(B), y \in Bx.$$

We expect  $x_\infty$  of (a) above to solve  $0 \in x_\infty + Bx_\infty$ , but cannot in fact state so without further assumptions on  $B$ . (As we will see, (b) is close to the assertion  $0 \in x_\infty + Bx_\infty$ ). Theorem 1 leaves open the question of existence of admissible sequences, which also requires further conditions on  $B$  (see Theorem 3 and Proposition 1 below). These questions are resolved for continuous  $B$  in the next result.

Theorem 2. Let  $B: D(B) \subseteq X \rightarrow X$  be continuous and accretive.

(a) Let  $D(B) = X$ . Given  $x_0 \in X$  inductively define

$$(1.3) \quad \begin{cases} h_{k+1} = 2^{-n_k}, & k = 0, 1, \dots \\ x_{k+1} = \frac{1}{1+h_{k+1}} x_k - \frac{h_{k+1}}{1+h_{k+1}} Bx_k, & k = 0, 1, \dots \end{cases}$$

where  $n_k$  is the least nonnegative integer  $n$  such that

$$(1.4) \quad \left\| B\left(\frac{2^n}{1+2^n} x_k - \frac{1}{1+2^n} Bx_k\right) - Bx_k \right\| < e^{-(h_1+h_2+\dots+h_k+1)}.$$

Then  $\{x_k\}$  is admissible for  $0 \in R(I + B)$ .

(b) Let  $\{x_k\}$  be admissible for  $0 \in R(I + B)$  and  $D(B)$  be closed. Then

$$x_k + Bx_k \rightarrow x_\infty + Bx_\infty = 0$$

where  $x_\infty = \lim_{k \rightarrow \infty} x_k$ .

Remarks: The scheme (1.3), (1.4) gives an algorithm for computing  $\{x_k\}$ . Our proofs will give explicit estimates on  $\|x_k - x_\infty\|$ , but we will not be able to estimate  $n_k$  (equivalently,  $h_k$ ) in general and therefore the number of steps required to achieve a preassigned accuracy. There are many alternatives to (1.3), (1.4). In particular, we mention that  $e^{-(h_1 + h_2 + \dots + h_k + 1)}$  on the right of (1.4) can be replaced by  $\psi(h_1 + \dots + h_k)$  for any decreasing integrable function  $\psi : (0, \infty) \rightarrow (0, \infty)$ .

If  $B$  is not necessarily continuous, we will use the condition

$$(R) \quad \inf_{\lambda > 0} \frac{d(R(I + \lambda(I + B)), x)}{\min(1, \lambda)} = 0 \quad \text{for } x \in \overline{D(B)},$$

where  $d(C, x)$  denotes the distance from  $C \subseteq X$  to  $x \in X$ .

Theorem 3. Let  $B$  be accretive and (R) hold. Then:

- (a) There is an admissible sequence  $\{x_k\}$  for  $0 \in R(I + B)$ .
- (b)  $0 \in \overline{R(I + B)}$  (the closure of  $R(I + B)$ ). In particular, if  $B$  is also closed, there is a solution  $x_\infty$  of  $0 \in x_\infty + Bx_\infty$  to which every admissible sequence converges.

Remarks: Suppose we knew that there is a subset  $D_0$  of  $\overline{D(B)}$  with the following property: For every  $\varepsilon > 0$  and  $x_0 \in D_0$  there are finite sequences  $\{x_1, x_2, \dots, x_N\} \subseteq D_0$ ,  $\{h_1, \dots, h_N\} \subset (0, \infty)$  and  $y_1 \in Bx_1$  such that  $2 \geq \sum_{j=1}^N h_j \geq 1$  and  $\sum_{i=0}^N \|x_{i+1} - x_i + h_{i+1}(x_{i+1} + y_{i+1})\| < \varepsilon$ . Then there will be admissible sequences for  $0 \in R(I + B)$ , as is easy to see. Various conditions which imply this property are discussed by M. Pierre [7,8]. We have singled out (R) here because of its simplicity and the fact that it falls



outside the scope of previous works. In particular, the requirement that  $\lim_{\lambda \downarrow 0} \inf \lambda^{-1} d(R(I + \lambda(I+B)), x) = 0$  for  $x \in \overline{D(B)}$  (which allows use of the results of Y. Kobayashi [3]) implies (R) but is not implied by it. However, none of the above mentioned conditions are necessary for the existence of admissible sequences. We note:

Proposition 1. Let  $B$  be accretive and  $0 \in \overline{R(I+B)}$ . Then there is an admissible sequence for  $0 \in R(I+B)$ . Conversely, if there is an admissible sequence for  $0 \in R(I+B)$  then  $0$  is in the closed convex hull of  $R(I+B)$ .

Remarks: (i) For exposition's sake we have discussed the problem  $0 \in R(I+B)$  where  $B$  is accretive. All our results adapt at once to the case  $0 \in R(A)$  when  $A - \omega I$  is accretive for some  $\omega > 0$ .

(ii) It would be nice to know if the existence of an admissible sequence for  $0 \in R(I+B)$  implies  $0 \in \overline{R(I+B)}$ .

## Section 2.

We begin with three lemmas which contain the heart of the proofs of Theorem 1-3. Lemma 1 and 2 (when  $A = 0$ ) are variations of estimates used by Kobayashi [3], (see also [2]). Theorems 1 and 3 are proved following the lemmas. The proof of Theorem 2, which is closely related to that of Theorem 3, is given next. The section ends with the proof of Proposition 1.

Hereafter  $B$  is accretive,  $x_k \in X$ ,  $k = 0, 1, \dots$  and  $y_k \in Bx_k$ ,  $k = 1, 2, \dots$  are regarded as given. Let  $\{h_k\}_{k=1}^{\infty}$  be a sequence of real positive numbers and define  $z_k \in X$  for  $k = 1, 2, \dots$  by

$$(2.1) \quad x_{k+1} - x_k + h_{k+1}(x_{k+1} + y_{k+1}) = h_{k+1}z_{k+1} \quad k = 0, 1, 2, \dots$$

Lemma 1. Let  $i \geq \ell > 0$  and (2.1) hold. Then

$$(2.2) \quad \|x_i - x_\ell\| \leq b_{i,\ell} \|x_\ell + y_\ell\| + \sum_{k=\ell+1}^i h_k \|z_k\|$$

where  $b_{i,\ell} = \min(1, \sum_{k=\ell+1}^i h_k)$ .

Proof: From (2.1) we have

$$\begin{aligned} h_{k+1} \|z_{k+1}\| + h_{k+1} \|x_\ell + y_\ell\| &\geq \|(1 + h_{k+1})(x_{k+1} - x_\ell) - (x_k - x_\ell) \\ + h_{k+1}(y_{k+1} - y_\ell)\| &\geq \|(1 + h_{k+1})(x_{k+1} - x_\ell) + h_{k+1}(y_{k+1} - y_\ell)\| \\ &\quad - \|x_k - x_\ell\| \geq (1 + h_{k+1}) \|x_{k+1} - x_\ell\| - \|x_k - x_\ell\| \end{aligned}$$

where the first and second estimates are the triangle inequality and the third follows from the accretiveness of  $B$ . Thus

$$(2.3) \quad \|x_{k+1} - x_\ell\| \leq (1 + h_{k+1})^{-1} \|x_k - x_\ell\| + h_{k+1} (1 + h_{k+1})^{-1} (\|x_\ell + y_\ell\| + \|z_{k+1}\|),$$

and iterating this from  $k = \ell$  to  $k = i-1$ ,

$$\begin{aligned} (2.4) \quad \|x_i - x_\ell\| &\leq \sum_{j=\ell+1}^i \prod_{m=j}^i (1 + h_m)^{-1} h_j (\|x_\ell + y_\ell\| + \|z_j\|) \\ &\leq M_{i,\ell} \|x_\ell + y_\ell\| + \sum_{j=\ell+1}^i h_j \|z_j\|, \end{aligned}$$

where

$$(2.5) \quad M_{i,\ell} = \sum_{j=\ell+1}^i \prod_{m=j}^i (1 + h_m)^{-1} h_j \leq \sum_{j=\ell+1}^i h_j.$$

Now

$$(2.6) \quad M_{i,\ell} \text{ is nondecreasing in } i \geq \ell+1 \text{ and } 0 \leq M_{i,\ell} \leq 1$$

since  $M_{i+1,\ell} = \frac{1}{(1+h_{i+1})} M_{i,\ell} + \frac{h_{i+1}}{1+h_{i+1}}$  is a convex combination of 1

and  $M_{i,\ell}$ . Thus (2.2) follows from (2.4) because  $M_{i,\ell} \leq b_{i,\ell}$  by (2.5) and (2.6).

**Lemma 2.** Let (2.1) be satisfied,  $A, B \geq 0$ ,  $\ell > 0$  and assume that

$$\|x_i - x_\ell\| \leq A + B \sum_{k=\ell+1}^i h_k + \sum_{k=\ell+1}^i h_k \|z_k\|$$

for  $i \geq \ell$ . Then for  $i \geq j \geq \ell$

$$(2.7) \quad \|x_i - x_j\| \leq A \prod_{k=\ell+1}^j (1 + h_k)^{-1} + B \sum_{k=j+1}^i h_k + \sum_{k=\ell+1}^i h_k \|z_k\| + \sum_{k=\ell+1}^j h_k \|z_k\|.$$

**Proof:** Set  $a_{i,j} = \|x_i - x_j\|$ . From the accretiveness of  $B$  and (2.1) we obtain (see, e.g. [2] Lemma 1.7 and Remark 4.1) the following recursion relation

$$(2.8) \quad a_{i,j} \leq \frac{h_i + h_j}{h_i + h_j + h_i h_j} \left( \frac{h_j}{h_i + h_j} (a_{i-1,j} + h_i \|z_i\|) + \frac{h_i}{h_i + h_j} (a_{i,j-1} + h_j \|z_j\|) \right).$$

By assumption, (2.7) holds if  $j = \ell$  and clearly it holds if  $i = j$ . Thus it suffices to show that if  $i > j \geq \ell$  and the desired estimate (2.7) holds for  $a_{i-1,j}$  and  $a_{i,j-1}$  then it holds for  $a_{i,j}$ . However, this follows readily from (2.8).

**Lemma 3.** Let (2.1) hold and  $\sum_{k=1}^{\infty} h_k \|z_k\| < \infty$ . Then  $\lim_{k \rightarrow \infty} x_k$  exists.

**Proof:** There are two cases. First we assume  $\sum_{k=0}^{\infty} h_k = \infty$ . By Lemma 1

$$\|x_i - x_\ell\| \leq \|x_\ell + y_\ell\| + \sum_{k=\ell+1}^i h_k \|z_k\| \quad \text{for } i \geq \ell,$$

so by Lemma 2 (with  $A = \|x_\ell + y_\ell\|$ ,  $B = 0$ )



$$(2.9) \quad \|x_i - x_j\| \leq \|x_\ell + y_\ell\| \prod_{k=\ell+1}^j (1 + h_k)^{-1} + \sum_{k=\ell+1}^i h_k \|z_k\| + \sum_{k=\ell+1}^j h_k \|z_k\|$$

for  $i \geq j \geq \ell$ . Since  $\sum_{k=1}^{\infty} h_k = \infty$ ,  $\prod_{k=\ell+1}^{\infty} (1 + h_k)^{-1} = 0$ .

Thus by (2.9)

$$(2.10) \quad \limsup_{i,j \rightarrow \infty} \|x_i - x_j\| \leq 2 \sum_{k=\ell+1}^{\infty} h_k \|z_k\|$$

for all  $\ell$ . Letting  $\ell \rightarrow \infty$  we find that  $\{x_k\}$  is a Cauchy sequence and hence convergent.

The other possibility is that  $\sum_{k=0}^{\infty} h_k < \infty$ . This time we use the assertion

$$\|x_i - x_\ell\| \leq \|x_\ell + y_\ell\| \sum_{k=\ell+1}^i h_k + \sum_{k=\ell+1}^i h_k \|z_k\| \quad i \geq \ell,$$

from Lemma 1. Now Lemma 2 (with  $A = 0$ ,  $B = \|x_\ell + y_\ell\|$ ) implies

$$\|x_i - x_j\| \leq \|x_\ell + y_\ell\| \sum_{k=j+1}^i h_k + \sum_{k=\ell+1}^i h_k \|z_k\| + \sum_{k=\ell+1}^j h_k \|z_k\|.$$

Letting  $i, j \rightarrow \infty$  we again find (2.10) (since  $\sum h_k < \infty$ ), and  $\{x_k\}$  is Cauchy as before.

**Proof of Theorem 1.** Part (a) is an immediate consequence of Lemma 3. To

obtain part (b) let  $\lim_{k \rightarrow \infty} x_k = x_\infty$  and  $y \in Bx$ . Rewriting (2.1) we have

$$(2.11) \quad y_{k+1} - y = h_{k+1}^{-1} (x_k - x_{k+1} + h_{k+1}(z_{k+1} - x_{k+1} - y)).$$

Since  $B$  is accretive

$$(2.12) \quad \|x_{k+1} - x\| \leq \|x_{k+1} - x + \lambda(y_{k+1} - y)\| \quad \text{for } \lambda > 0.$$

Substitute (2.11) in (2.12), multiply by  $h_{k+1}/\lambda$  and manipulate to find:

$$\begin{aligned}
(1 + h_{k+1}) \|x_{k+1} - x\| &\leq \|x_k - x\| + h_{k+1} \|z_{k+1}\| + \left\| \left(1 - \frac{h_{k+1}}{\lambda}\right)(x - x_{k+1}) - h_{k+1}x_{k+1} - h_{k+1}y \right\| \\
&\quad - \left(\frac{h_{k+1}}{\lambda} - 1 - h_{k+1}\right) \|x_{k+1} - x\| \leq \|x_k - x\| + h_{k+1} \|z_{k+1}\| \\
&\quad + \left\| \left(1 - \frac{h_{k+1}}{\lambda}\right) - 1 - h_{k+1}\right\| (x - x_{k+1}) - h_{k+1}x_{k+1} - h_{k+1}y \left\| - \frac{h_{k+1}}{\lambda} \|x_{k+1} - x\|
\end{aligned}$$

and therefore

$$\begin{aligned}
(2.13) \quad (1 + h_{k+1}) \|x - x_{k+1}\| &\leq \|x - x_k\| + \lambda^{-1} h_{k+1} (\|x - x_{k+1} + \lambda(x+y)\| \\
&\quad - \|x - x_{k+1}\|) + h_{k+1} \|z_{k+1}\|.
\end{aligned}$$

Iterating this inequality from  $k = \ell$  to  $k = i-1$  yields

$$\begin{aligned}
(2.14) \quad \|x - x_i\| &\leq \|x - x_\ell\| \prod_{j=\ell+1}^i (1 + h_j)^{-1} + \\
&\quad \lambda^{-1} \sum_{j=\ell+1}^i \left( \prod_{m=j}^i (1 + h_m)^{-1} \right) h_j (\|x - x_j + \lambda(x+y)\| - \|x - x_j\|) \\
&\quad + \sum_{j=\ell+1}^i \left( \prod_{m=j}^i (1 + h_m)^{-1} \right) h_j \|z_j\|.
\end{aligned}$$

Since  $\sum_{j=1}^{\infty} h_j = \infty$ , the first term on the right of (2.14) tends to zero as  $i \rightarrow \infty$ .

The third term on the right also tends to zero as  $i \rightarrow \infty$  by the dominated convergence theorem (each term individually tends to 0 and  $\{h_j \|z_j\|\}$  is a dominating summable sequence). Finally, by (2.6) and

$$\lim_{j \rightarrow \infty} (\|x - x_j + \lambda(x+y)\| - \|x - x_j\|) = \|x - x_\infty + \lambda(x+y)\| - \|x - x_\infty\|$$

we obtain upon letting  $i \rightarrow \infty$  in (2.14) that

$$\|x - x_\infty\| \leq \lambda^{-1} (\|x - x_\infty + \lambda(x+y)\| - \|x - x_\infty\|)$$

or, after rearranging,

$$(1 + \lambda) \|x - x_\infty\| \leq \|x - x_\infty + \lambda(x+y)\|$$

which is the desired inequality.

Proof of Theorem 3. We begin by showing that if  $B$  is accretive and satisfies (R) then there is an admissible sequence for  $0 \in R(I + B)$ .

For  $x \in \overline{D(B)}$  and  $\varepsilon > 0$  let  $\Lambda(x, \varepsilon)$  be the set of those numbers  $\lambda > 0$  for which there exists  $x_\lambda$  and  $y_\lambda \in Bx_\lambda$  such that

$$\|x_\lambda + \lambda(x_\lambda + y_\lambda) - x\| < \chi(\lambda)\varepsilon$$

where  $\chi(\lambda) = \min(1, \lambda)$  for  $\lambda \in (0, \infty]$ .  $\Lambda(x, \varepsilon)$  is nonempty by condition (R),

and we define  $\lambda(x, \varepsilon) = \sup \Lambda(x, \varepsilon)$ . Let  $x_0 \in \overline{D(B)}$  be arbitrary and suppose

$x_1, x_2, \dots, x_{k-1}$  and  $h_1, h_2, \dots, h_{k-1}$  have been chosen. Set

$$(2.15) \quad \varepsilon_k = \exp\left(-\sum_{j=1}^{k-1} h_j - 1\right)$$

and choose  $h_k > 0$ ,  $x_k, y_k \in Bx_k$  so that

$$(2.16) \quad \begin{cases} \chi(\frac{1}{2}\lambda(x_{k-1}, \varepsilon_k)) \leq h_k < \infty \\ \|x_k - x_{k-1} + h_k(x_k + y_k)\| < \chi(h_k)\varepsilon_k. \end{cases}$$

In this way we get infinite sequences  $\{h_k\}$ ,  $\{x_k\}$ ,  $\{y_k\}$ . Now by (2.16),

(2.15) and with  $z_k$  as in (2.1),

$$(2.17) \quad \sum_{k=1}^{\infty} h_k \|z_k\| < \sum_{k=1}^{\infty} \chi(h_k)\varepsilon_k \leq \int_0^{\infty} e^{-s} ds < \infty.$$

Thus, if  $\sum_{k=1}^{\infty} h_k = \infty$ ,  $\{x_k\}$  is admissible and we are done. Let  $\sigma_k = h_1 + \dots$

$+ h_k$ . Assuming  $\lim_{k \rightarrow \infty} \sigma_k = \sigma_\infty < \infty$  we will reach a contradiction by use of a

now standard idea of Nagumo [6], and thus complete the proof of (a). By

Lemma 3,  $x_\infty = \lim_{k \rightarrow \infty} x_k$  exists. By (R) there exists  $\lambda_0 \in (0, \infty)$ ,  $\bar{x}, \bar{y} \in B\bar{x}$

such that

$$\|\bar{x} - x_\infty + \lambda_0(\bar{y} + \bar{x})\| < e^{-(\sigma_\infty + 1)} \chi(\lambda_0).$$



Then

$$\|\bar{x} - x_{k-1} + \lambda_0(\bar{y} + \bar{x})\| < e^{-(\sigma_{k-1}+1)} \chi(\lambda_0) = \chi(\lambda_0) \varepsilon_k.$$

for all  $k$  large enough. Hence, by (2.16),  $2h_k \geq \chi(\lambda_0)$  for large  $k$ , contradicting  $\sigma_k = h_1 + \dots + h_k \rightarrow \sigma_\infty < \infty$ .

To prove part (b), we use (R) to assert the existence of sequences

$\lambda_i > 0$ ,  $u_i \in D(B)$ ,  $v_i \in Bu_i$  such that

$$(2.18) \quad \left\| \frac{u_i - x_\infty}{\lambda_i} + (u_i + v_i) \right\| \rightarrow 0.$$

By Theorem 1(b) we also have

$$(2.19) \quad (1 + \lambda_i) \left\| \frac{u_i - x_\infty}{\lambda_i} \right\| \leq \left\| \frac{u_i - x_\infty}{\lambda_i} + (u_i + v_i) \right\|.$$

Combining (2.18) and (2.19) we conclude  $\lambda_i^{-1}(u_i - x_\infty) \rightarrow 0$  as well as  $u_i \rightarrow x_\infty$  and subsequently, from (2.18),  $u_i + v_i \rightarrow 0$ . Thus  $0 \in \overline{R(I+B)}$  and  $0 \in R(I+B)$  if  $B$  is closed.

Proof of Theorem 2. We first show that (4), (5) defines an admissible sequence. With  $y_k = Bx_k$  we have, by (4), (5)

$$\begin{aligned} \sum_{k=0}^{\infty} \|x_{k+1} - x_k + h_{k+1}(x_{k+1} + y_{k+1})\| &= \sum_{k=0}^{\infty} h_{k+1} \|Bx_{k+1} - Bx_k\| \\ &\leq \sum_{k=0}^{\infty} h_{k+1} \exp\left(-\sum_{j=1}^k h_j - 1\right) \leq 1. \end{aligned}$$

Thus it is sufficient to show  $\sum_{k=0}^{\infty} h_j = \infty$ . Suppose  $\sigma_k = h_1 + \dots + h_k \rightarrow \sigma_\infty < \infty$ . Then  $x_k$  converges to a limit  $x_\infty$  by Lemma 3. Since  $B$  is continuous there is an integer  $n$  such that

$$\left\| B\left(\frac{2^n}{1+2^n} x_\infty - \frac{1}{1+2^n} Bx_\infty\right) - Bx_\infty \right\| < \exp(-\sigma_\infty - 1).$$

Then also

$$\|B(\frac{2^n}{1+2^n}x_k - \frac{1}{1+2^n}Bx_k) - Bx_k\| < \exp(-(h_1 + \dots + h_k)-1)$$

for large  $k$  and therefore  $n_k \leq n$  for large  $k$ . But then  $\sigma_k = h_1 + \dots + h_k \rightarrow \infty$  by (4), a contradiction.

Let  $\{x_k\}$  be an admissible sequence for  $0 \in R(I+B)$  and (2.1) hold with  $\sum_{k=1}^{\infty} h_k \|z_k\| < \infty$ . Then  $x_k \rightarrow x_{\infty}$  and summing (2.1) from  $k=m$  to  $k=n-1$  yields

$$\sum_{j=m+1}^n \frac{h_j}{\sigma_n - \sigma_m} (x_j + y_j) = \frac{x_m - x_n}{\sigma_n - \sigma_m} + \frac{1}{\sigma_n - \sigma_m} \sum_{j=m+1}^n h_j z_j,$$

where  $\sigma_j = h_1 + \dots + h_j$ . The right hand side above tends to zero as  $n, m \rightarrow \infty$  subject to  $\sigma_n - \sigma_m \geq 1$ , while the left hand side consists of convex combinations of  $x_j + y_j$ . Thus if  $x_j + y_j$  has a limit as  $j \rightarrow \infty$ , it must be zero.

If  $B$  is continuous and  $D(B)$  is closed,  $x_{\infty} \in D(B)$  and  $x_j + Bx_j \rightarrow x_{\infty} + Bx_{\infty}$ , proving (b).

Proof of Proposition 1. Let  $0 \in \overline{R(I+B)}$ . Let  $\{\alpha_k\}$  be an arbitrary summable sequence of positive numbers and  $\{h_k\}$  a sequence of positive numbers satisfying  $\sum_{k=1}^{\infty} h_k = \infty$ ,  $\sum_{k=1}^{\infty} \alpha_k h_k < \infty$ . Choose  $x_k \in D(B)$ ,  $y_k \in Bx_k$  such that  $\|x_k + y_k\| < \alpha_k$ . Then we claim  $\{x_k\}$  is an admissible sequence for  $0 \in R(I+B)$  and we may use  $\{h_k\}$  in (3). Indeed

$$\begin{aligned} \|x_{k+1} - x_k\| &\leq \|x_{k+1} - x_k + y_{k+1} - y_k\| \leq \|x_{k+1} + y_{k+1}\| + \|x_k + y_k\| \\ &\leq \alpha_{k+1} + \alpha_k \end{aligned}$$

$$\begin{aligned} \text{so} \\ \sum_{k=1}^{\infty} \|x_{k+1} - x_k + h_{k+1}(x_{k+1} + y_{k+1})\| &\leq \sum_{k=1}^{\infty} \|x_{k+1} - x_k\| + h_{k+1} \|x_{k+1} + y_{k+1}\| \\ &\leq 2 \sum_{k=1}^{\infty} \alpha_k + \sum_{k=1}^{\infty} \alpha_k h_k < \infty. \end{aligned}$$

Thus  $\{x_k\}$  is admissible.

In the proof of Theorem 2 we showed that certain convex combinations of  $x_j + y_j$  converged to 0 if (3) holds, so 0 is in the closed convex hull of  $R(I + B)$  if there is an admissible sequence. This completes the proof.

Remarks: From the proof of proposition 1 we see that if  $0 \in \overline{R(I + B)}$  then there exist admissible sequences with arbitrary associated sequences  $\{h_k\}$  satisfying  $\sum h_k = \infty$ . It is worth noting that if (3) holds and  $\inf_k h_k > 0$  then  $0 \in \overline{R(I + B)}$  for

$$h_k(x_k + y_k) = (x_{k-1} - x_k + h_{k+1}(x_{k+1} + y_{k+1})) - (x_{k-1} - x_k)$$

and the first term on the right tends to 0 by (3) (ii) and the second does by

Theorem 1, so  $x_k + y_k \rightarrow 0$ . One can also show that  $x_k + y_k \rightarrow 0$  if

$$\sum_{k=1}^{\infty} \|z_k\| < \infty.$$

### 3. Applications.

In this section we deduce two known general theorems on accretive sets in Banach space as simple consequences of our results. We note that the previously known proofs of these theorems were all dependent on the existence of a solution to the initial value problem (1).

We start by introducing the following conditions:

$$(R_1) \quad \liminf_{\lambda \rightarrow 0^+} \lambda^{-1} \text{dist}(R(I + \lambda A); x - \lambda y) = 0 \quad \forall x \in \overline{D(A)}, y \in X.$$

Note that it is sufficient for  $(R_1)$  to hold only for every  $y$  in a dense subset of  $X$  in order for it to hold for every  $y \in X$ .

#### Lemma 4.

Let  $A$  be accretive and satisfy  $(R_1)$ . If  $P: \overline{D(A)} \subset X \rightarrow X$  is continuous, then  $A+P$  satisfies  $(R_1)$ .



Proof:

If  $A$  satisfies  $(R_1)$  then for every  $y \in X$  there are sequences  $\lambda_i \rightarrow 0$  and  $y_i \in Ax_i$  such that

$$(3.1) \quad \lambda_i^{-1}(x_i + \lambda_i y_i - (x - \lambda_i y - \lambda_i Px)) \rightarrow 0 \quad \text{as } i \rightarrow \infty.$$

But

$$\begin{aligned} \lambda_i^{-1}(x_i + \lambda_i y_i + \lambda_i Px_i - (x - \lambda_i y)) &= \lambda_i^{-1}(x_i + \lambda_i y_i - (x - \lambda_i y - \lambda_i Px)) \\ &\quad + Px_i - Px. \end{aligned}$$

In order to prove that  $A+P$  satisfies  $(R_1)$  it is therefore sufficient to show that  $x_i \rightarrow x$  as  $i \rightarrow \infty$ . From the accretiveness of  $A$  we have

$$\|x_i - u\| \leq \|x_i - u + \lambda_i(y_i - v)\| \quad \text{for } v \in Au$$

and from (3.1) we then have

$$\limsup_{i \rightarrow \infty} \|x_i - u\| \leq \limsup_{i \rightarrow \infty} \|x - u - \lambda_i(y + Px + v)\| = \|x - u\|$$

for all  $u \in D(A)$  and thus  $x_i \rightarrow x$ .

Let  $z \in X$  be arbitrary. Taking  $P = I - z$  in the previous lemma, it follows that  $A_z = A + I - z$  satisfies condition  $(R_1)$  and hence also  $(R)$  of Section 1 and therefore by Theorem 3 we have:

Theorem (Y. Kobayashi [3]).

Let  $A$  be accretive and satisfy  $(R_1)$ . Then  $\bar{A}$  (the closure of  $A$ ) is  $m$ -accretive.

We conclude with the following general perturbation theorem.

Theorem (Y. Kobayashi [3]).

Let  $A$  be accretive and  $P : \overline{D(A)} \rightarrow X$  be continuous. If  $A + P$  is accretive, then it is  $m$ -accretive if and only if  $A$  is  $m$ -accretive.

Proof:

Let  $A$  be  $m$ -accretive then  $A$  is closed and satisfies  $(R_1)$ . Therefore, by Lemma 4,  $A+P$  satisfies  $(R_1)$ . Since, by the continuity of  $P$ ,  $A+P$  is closed and by assumption it is accretive, it follows from the previous theorem that  $A+P$  is  $m$ -accretive. If  $A+P$  is  $m$ -accretive one has that  $A = (A+P)-P$  is as well by the above.

The last theorem is a considerable generalization of the theorem of R. Martin which was stated in the introduction. Indeed Martin's theorem is obtained by taking  $A \equiv 0$  on  $X$ . G. Webb [9] proved the above perturbation result assuming that  $A$  was linear  $m$ -accretive with  $\overline{D(A)} = X$ . Subsequently V. Barbu [2] generalized Webb's result to the case where  $A$  was a general  $m$ -accretive operator and  $P : X \rightarrow X$  was continuous. Finally Y. Kobayashi [3] proved the above theorem.

References

- [1] V. Barbu, Continuous perturbations of nonlinear  $m$ -accretive operators in Banach space, Boll. d. U. Mat. Ita. 6 (1972), pp. 270-278.
- [2] M. G. Crandall and L. C. Evans, On the relation of the operator  $\frac{\partial}{\partial \tau} + \frac{\partial}{\partial s}$  to evolution governed by accretive operators, Israel J. Math. 21 (1975), 261-278.
- [3] Y. Kobayashi, Difference approximation of Cauchy problems for quasi-dissipative operators and generation of nonlinear semigroups, J. Math. Soc. Japan 27 (1975), 640-665.
- [4] R. H. Martin, A global existence theorem for autonomous differential equations in Banach space, Proc. Amer. Math. Soc. 26 (1970), 307-314.

- [5] G. H. Minty, Monotone (nonlinear) operators in a Hilbert space,  
Duke Math. J. 29 (1962), 341-346.
- [6] M. Nagumo, Über die lage der Integralkurven gewöhnlicher  
Differentialgleichungen, Proc. Phys. - Math. Soc. Japan 24 (1942),  
551-559.
- [7] M. Pierre, Un théorème général de génération de semi-groupes non-  
linéaires, Israel J. Math. , 23 (1976), 189-199.
- [8] M. Pierre, Génération et perturbation de semi-groupes de contractions  
non lineaires, These de Docteur de 3<sup>e</sup> cycle, Universite de Paris VI,  
1976.
- [9] G. F. Webb, Nonlinear perturbations of linear accretive operators in  
Banach spaces, J. Func. Anal. 10 (1972), 191-203.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1682 (14) MRQ-TSR-1682	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) On the Range of Accretive Operators. ✓	5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period	
7. AUTHOR(s) (10) Michael G./Crandall Amnon/Pazy	8. CONTRACT OR GRANT NUMBER(s) (15) DAAG29-75-C-0024, ✓ NSF-MCS-76-10227	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (12) 19 p.	
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below	12. REPORT DATE (11) Sep 1976	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 16	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) (9) Technical summary rept.		
18. SUPPLEMENTARY NOTES U. S. Army Research Office and National Science Foundation P.O. Box 12211 Washington, D. A. 20550 Research Triangle Park, N.C. 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Accretive operator, iteration scheme, range		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) → The solvability of the nonlinear operator equation $w = x + Bx$ , where $B$ is accretive in a general Banach space $X$ is studied by means of discrete approximations. In particular, an algorithm is given for solving the equation in the case that $B$ is continuous and everywhere defined. ←		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED 221 200

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)